

## YOUR CURRENT MEDICATION INSTRUCTIONS

## DO NOT TAKE MEDICATIONS THAT CAN CAUSE BLEEDING

DO NOT TAKE ANY ASPIRIN, IBUPROFEN AND RELATED MEDICATIONS AT LEAST 2 WEEKS PRIOR TO SURGERY! If YOU DO NOT STOP TAKING, WE WILL CANCEL YOUR SURGERY AND YOU WILL LOSE YOUR DEPOSIT!

## Common Medications that can thin your blood TO STAY AWAY FROM 2 WEEKS PRIOR TO SURGERY:

- ·ASPIRIN (Excedrin)
- •Oil Based Vitamins and Supplements- Fish/ Krill Oil, Vitamin A/D/E, and Garlic Tablets.
- ·Blood thinners- Coumadin/Warfarin, Plavix, Xarelto
- **·IBUPROFEN**
- ·Naxoproxen (Naprosyn, Aleve)

Tylenol is the only non-prescription pain medication that is OK to take before surgery.

If you have taken aspirin please notify the office immediately. Your surgery may be cancelled.

PLEASE DISCUSS ALL CURRENT MEDICATIONS WITH YOUR SURGEON TO DETERMINE IF THEY NEED TO BE STOPPED.

\*\*\*YOU WILL NEED TO STOP ALL VITAMINS, AND HERBAL SUPPLEMENTS FOR 7 DAYS. THE DAY BEFORE SURGERY AND THE DAY OF SURGERY, you may continue them 7 days after your surgery \*\*\*

Name (print)	Signature	Date	
 Staff Name (print)	Signature	Date	